



Cell name: A-431 x ABCG2

Cat. No. A-0340

Product description


A-431 x ABCG2 was established by retroviral transduction of human ATP-binding cassette, sub-family G, member 2 cDNA into A-431 cells and flow cytometry sorting of the anti-Human ABCG2 (clone 5D3) immunolabeled population.

Transgene construct information

Expression cassette type

Single-promoter, monocistronic

Expression cassette map



Expression cassette features

Element	Type	Species	RefSeq	Mutation / Discrepancy
SFFV	promoter, viral, constitutive	-	-	-
ABCG2	CDS	Homo sapiens	NM_004827.2	none / none

Transgene protein information

The transmembrane protein encoded by this gene is included in the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the White subfamily. Alternatively referred to as a breast cancer resistance protein, this protein functions as a xenobiotic transporter which may play a major role in multi-drug resistance. It likely serves as a cellular defense mechanism in response to mitoxantrone and anthracycline exposure. Significant expression of this protein has been observed in the placenta, which may suggest a potential role for this molecule in placenta tissue. Multiple transcript variants encoding different isoforms have been found for this gene. [Source: NCBI Gene, <http://www.ncbi.nlm.nih.gov/gene/9429>]

Cell culture characteristics

Source

Human skin epidermoid carcinoma

Morphology

Epithelial

Growth properties

Adherent

Culture conditions

Culture cells at 37°C in humidified atmosphere with 5% CO₂. The base medium for this cell line is α-MEM with 4.5 g/l glucose. To make the complete growth medium, add fetal bovine serum to a final concentration of 10%.

Subculturing

Remove culture medium, rinse twice thoroughly with PBS, and detach cells by treating with 0.04 mL/cm² of 0.25% trypsin / 1 mM EDTA solution for 15 minutes at 37°C. Add 5x volume of complete medium to neutralize trypsin-EDTA, pellet cells at 250 x g, discard supernatant, and resuspend in culture medium. Subcultivate in a ratio of 1:4-1:6 three times a week. Recommended plating density: 3x10⁴ cells/cm².

Preservation

Freeze Medium: Complete growth medium with 10% DMSO.

Storage Temperature: Liquid nitrogen vapour phase.

Population doubling time

~30 hours


Sterility testing

Mycoplasma: negative

Comment

This cell line overexpresses Epidermal Growth Factor Receptor (EGFR).

Validation results



A: Red line: A-431 x ABCG2 cells incubated with anti-Human ABCG2 antibody.

Dotted line: A-431 x ABCG2 cells incubated with isotype control.

Indirect immunofluorescent labeling with anti-Human ABCG2 (clone 5D3) antibody (R&D Systems®) or isotype control + AlexaFluor™ 488 labeled anti-Mouse IgG secondary antibody. Detection by flow cytometry.

B: Red line: A-431 x ABCG2 cells incubated with mitoxantrone and Ko143.

Dotted line: A-431 x ABCG2 cells incubated with mitoxantrone.

Mitoxantrone uptake detected by flow cytometry at 633 nm in the presence or absence of Ko143, a specific inhibitor of ABCG2.