

Cell name: HCC827 x ABCB1-Neo
Cat. No. HC-0220

Product description


HCC827 x ABCB1-Neo was established by retroviral transduction of human ABCB1 (also known as P-glycoprotein or MDR1) cDNA into HCC827 cells and flow cytometry sorting of the calcein-extruding population.

Transgene construct information

Expression cassette type

Single-promoter, bicistronic

Expression cassette map



Expression cassette features

Element	Type	Species	RefSeq	Mutation / Discrepancy
SFFV	promoter, viral, constitutive	-	-	-
Splice donor site	Splice donor site	-	-	-
ABCB1	CDS	Homo sapiens	AF016535	none / none
Splice acceptor site	Splice acceptor site	-	-	-
Neomycin resistance gene	CDS	-	-	none / none

Transgene protein information

The transmembrane protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the MDR/TAP subfamily. Members of the MDR/TAP subfamily are involved in multidrug resistance. The protein encoded by this gene is an ATP-dependent drug efflux pump for xenobiotic compounds with broad substrate specificity. It is responsible for decreased drug accumulation in multidrug-resistant cells and often mediates the development of resistance to anticancer drugs. This protein also functions as a transporter in the blood-brain barrier. [Source: NCBI Gene, <http://www.ncbi.nlm.nih.gov/gene/5243>]

Creative Cell Ltd.

Homepage: www.creativecell.hu Email: info@creativecell.hu Phone: +36 (30) 41-44446

Cell culture characteristics

Source

Human lung adenocarcinoma

Morphology

Epithelial

Growth properties

Adherent

Culture conditions

Culture cells at 37°C in humidified atmosphere with 5% CO₂. The base medium for this cell line is RPMI-1640 with 4.5 g/l glucose. To make the complete growth medium, add fetal bovine serum to a final concentration of 10%.

Subculturing

Remove culture medium and detach cells by treating with 0.04 mL/cm² of 0.25% trypsin / 1 mM EDTA solution for 10 minutes at 37°C. Add 5x volume of complete medium to neutralize trypsin-EDTA, pellet cells at 250 x g, discard supernatant, and resuspend in culture medium. Subcultivate in a ratio of 1:4-1:6 three times a week. Recommended plating density: 4x10⁴ cells/cm².

Preservation

Freeze Medium: Complete growth medium with 10% DMSO.

Storage Temperature: Liquid nitrogen vapour phase.

Population doubling time

~30 hours


Sterility testing

Mycoplasma: negative

Comment

This lung adenocarcinoma cell line has an acquired mutation in the EGFR tyrosine kinase domain (E746 - A750 deletion).

Validation results



A: Red line: HCC827 x ABCB1-Neo cells incubated with anti-Human ABCB1 antibody.

Dotted line: HCC827 x ABCB1-Neo cells incubated with isotype control.

Indirect immunofluorescent labeling with anti-Human ABCB1 antibody (clone MRK16, Alexis Biochemicals / Enzo Life Sciences) or isotype control + PE labeled anti-Mouse IgG secondary antibody. Detection by flow cytometry.

B: Red line: HCC827 x ABCB1-Neo cells incubated with calcein.

Dotted line: HCC827 x ABCB1-Neo cells incubated with calcein and verapamil.

Calcein uptake was detected by flow cytometry at 485 nm in the presence or absence of verapamil, an inhibitor of ABCB1.

Creative Cell Ltd.

Homepage: www.creativecell.hu Email: info@creativecell.hu Phone: +36 (30) 41-44446